
Knowability and no ability in climate projections

Gerard Roe,

Earth Space Sciences, Dept. Washington, Seattle, WA



How sensitive is climate to changes in CO₂?

A traditional measure

• Climate sensitivity (or equilibrium climate sensitivity)


Definition: the long-term change in annual-mean, global-mean, near-surface air temperature to a doubling of CO₂ above preindustrial values (phew!, e.g., Arhenius, 1896, Charney, 1979)

• IPCC 2007 says:


Likely (2-in-3)
$$2.0 < \Delta T < 4.5$$
°C
Very unlikely (<1-in-10) $\Delta T < 1.5$ °C

• Note this leaves \sim 2-in-10 chance for Δ T > 4.5 °C (though IPCC says observations are less well fit with these values)


1. Different estimates




1. Different estimates



1. Different estimates




1. Different estimates



So why these values, and why this shape?

1.5 An aside



- The main IPCC climate models under-sample the allowed range.
- An issue for regional climate predictions?

2. Estimates from observations

Global energy budget:


$$R_f$$
 = F + $\lambda^{-1}\Delta 7$ forcing = storage + atmospheric response

In principle, get R_f , F, ΔT from observations, solve for λ , then:

$$\Delta T_{2xCO_2} = \lambda R_{f2xCO_2}$$


2. Estimates from observations

How much warming has there been since pre-industrial times?



• Global mean temperature change is <u>well observed</u>.


2. Estimates from observations



Numbers from IPCC, 2007

• Warming from CO₂ and other Greenhouse gases (CH₄, O₃) (plus a tiny bit from solar)

2. Estimates from observations




Numbers from IPCC, 2007 and Lyman et al. (2010)

Cooling from heat storage in ocean, and aerosols

Aerosols: airborne particulates (solid/liquid)


have complicated effects (some warm, some cool, change clouds)

2. Estimates from observations



 Total climate forcing is <u>quite uncertain</u> and aerosols are the culprit.


3. Estimates from observations



$$\lambda = \frac{\Delta T}{R_f - H}$$


• Fat tail is because aerosol forcing could be quite negative

3. Estimates from models




 Black curve is the relationship between climate feedbacks and climate sensitivity.

3. Estimates from models




Green curve reflects current uncertainty in climate feedbacks.

3. Estimates from models



Red curve is resulting uncertainty in climate sensitivity.

3. Estimates from models



Red curve is resulting uncertainty in climate sensitivity.

4. Prospects for progress

a. Improved observations/models

Its hard!! Incremental improvements, but probably no breakthroughs.

b. Combine different estimates?

Very hard to establish the <u>degree of independence</u> of individual estimates. (see Knutti and Hegerl, 2008)

c. Use other observations?

(e.g., NH vs. SH; pole-to-eq. ΔT ; seasonality, trop. water vapor)


Structural errors among models highly uncertain. (see Knutti et al, 2010)

→ Prudent not to expect big improvements any time soon....

1. What if all anthropogenic emissions ceased tomorrow?

Lifetimes: CO₂: centuries to 100,000 yrs+


Aerosols: days to weeks



1. What's already in store for us?

Lifetimes: CO₂: centuries to 100,000 yrs+


Aerosols: days to weeks



Immediate loss of aerosols unmasks GHG gas warming


1. What's already in store for us?

Idealized timeline of past and future climate forcing, if we stop everything today




1. What's already in store for us?

Our best guess at what would happen




1. What's already in store for us?

But if past forcing has been high....



1. What's already in store for us?

But if past forcing has been low....



2. Past forcing and climate sensitivity are intrinsically related

If past forcing is strong -> climate sensitivity is low.


If past forcing is weak → climate sensitivity is high.

For Integrated Assessment Models this matters:

 forcing (including aerosol forcing) cannot be assumed to be independent of climate sensitivity.

Transient evolution of climate

1. Heat uptake of the ocean is diffusive




Hansen et al. (1985) show this means that

Climate adjustment time is proportional to (Climate Sensitivity)²

Transient evolution of climate

2. The fat tail grows very slowly




climate model response (mean & 95% bounds) to an instantaneous doubling of CO₂

 Constraining the details of the far tail of climate sensitivity is not useful on societally relevant timescales?

CO₂ stabilization targets are a mistake

1. Climate response to fixed level of CO₂ is uncertain




(Allen and Frame, 2007)
Stabilization target
of 450 ppm at 2100

- High end sensitivities take a long, long time to be realized
- There is still considerable uncertainty at 2150.

CO₂ stabilization targets are a mistake

2. Flexibility is key




(Allen and Frame, 2007)

Concentration target adjusted at 2050.

A flexible emissions strategy is key to reaching a desired goal

Does global climate predict local climate?

- 1. Is climate sensitivity a good predictor of regional change?
- Among models, how well are varns in global climate sensitivity correlated with varns in regional climate change at 2100?



If |corr. coeff.|
< 0.70 then
<50% of local
change is
associated with
global mean
change.

- The magnitude of local changes is affected by many factors
- Global ΔT is quite a poor predictor of local ΔT , ΔP
- If impacts are local, should global ΔT be used to calculate damages?


19 models from IPCC 2007 report, For more calculations see my web site. (calcⁿs made by Nicole Feldl)

Summary:




1. Uncertainty is not ignorance.


The planet is warming and its us that's doing it.




2. Climate sensitivity is uncertain b/c past forcing is uncertain (primarily aerosols).




3. Uncertainty in climate sensitivity and climate forcing are not independent.



4. <u>If</u> climate sensitivity is high, it takes a very long time to get there.





5. CO₂ stabilization targets are not an efficient way to achieve a climate goal. (flexibility is vital)




6. Global climate is not a strong predictor of local climate change.

Extra slides....







AR4 models undersample climate commitment




- Dark blue is the IPCC 'likely' (68% confidence interval) range of climate sensitivity (2 to 4.5 C) and implied range of radiative forcing
- AR4 climate models span only this 'likely' range
- R and λ are correlated within AR4 and older models (Kiehl 2007, Knutti 2008)

Effects of nonlinearity of climate feedbacks






By how much do observations have to change to change climate sensitivity



Aspects of feedbacks III.

How does uncertainty in feedbacks translate into uncertainty in the system response?



Systems of strong positive feedbacks inherently less predictable